

D

1. Which of the following phenomena could not be explained by classical physics and required a quantum hypothesis in order to make theory agree with experiment?

- | | |
|---|-----------------------|
| (a) Blackbody spectra | (e) Friction |
| (b) Diffraction of light | (f) (a), (c), and (d) |
| (c) Low-temperature heat capacity in perfect crystals | (g) (a), (b), and (d) |
| (d) The photoelectric effect | (h) All of the above |

2. Which of the following did Bohr assume in order to derive a model consistent with the photoemission spectrum of the hydrogen atom?

- | | |
|--|--|
| (a) The electron orbits the nucleus like a planet does a sun | (e) The ionization potential is equal to the work function |
| (b) The angular momentum of the electron is quantized | (f) (a) and (b) |
| (c) The one-electron atom is like a particle in a box | (g) (b) and (d) |
| (d) The Coulomb potential is quantized | (h) None of the above |

3. Which of the following statements about the de Broglie wavelength λ are *true*?

- | | |
|---|---|
| (a) λ decreases as mass increases if velocity is constant | (e) A particle that has zero velocity has an infinite de Broglie wavelength |
| (b) $\lambda = h / p$ | (f) All of the above |
| (c) λ decreases as momentum increases | (g) (a), (b) and (e) |
| (d) λ increases as kinetic energy decreases | (h) (c) and (d) |

4. Which of the following statements about a well behaved wave function is *false*?

- | | |
|---|--|
| (a) It must be continuous | (e) Its square modulus has units of probability density |
| (b) It may take on complex values | (f) It must be an eigenfunction of the momentum operator |
| (c) It must be quadratically integrable | (g) (d) and (f) |
| (d) It must be equal to its complex conjugate | (h) (a), (b), (c), and (e) |

NAME: _____

5. Which of the below equations will be true for any arbitrary pair of orthonormal functions f and g ?

(a) $\langle |f|^2 \rangle \langle |g|^2 \rangle = 1$

(b) $\langle f | H | g \rangle = 0$

(c) $\langle f | g \rangle = 0$

(d) $fg = 0$

(e) $f^*g - g^*f = 0$

(f) (a) and (c)

(g) (b), (d) and (e)

(h) All of the above

6. Which of the below expectation values are zero?

(a) $\langle \sin x | x | \cos x \rangle$

(b) $\langle \sin^2 x | x | \cos^2 x \rangle$

(c) $\langle f | [A, B] | g \rangle$ where A and B commute

(d) $\langle \Psi | H | \Psi \rangle$ where Ψ is a stationary state

(e) $\langle f | g \rangle - \langle g | f \rangle$

(f) $\langle \mu_{mn} \rangle$ for a forbidden transition

(g) (b), (d) and (f)

(h) (b), (c), and (f)

7. If a normalized wave packet Ψ is given as $\Psi(x, y, z, t) = \sum_{n=1}^{\infty} c_n \psi_n(x, y, z) e^{-iE_n t / \hbar}$,

what is the probability that an experiment will cause the system to collapse to the specific stationary state j ?

(a) Quantum mechanics does not allow you to know this probability

(b) $\langle \psi_j | H | \psi_j \rangle$

(c) One

(d) $|c_j|^2$

(e) c_j

(f) c_j^*

(g) (b) and (d)

(h) Only Schrödinger's cat knows

NAME: _____

8. Which of the following statements are *false* about the free particle?

- | | |
|---|--|
| <p>(a) Its Schrödinger equation is</p> $\left(-\frac{\hbar^2}{2m} \frac{d^2}{dx^2} - E\right)\Psi(x) = 0$ | <p>(e) Valid wave functions include</p> $\Psi(x) = Ae^{ikx} + Be^{-ikx}$ <p>where</p> $k = \frac{\sqrt{2mE}}{\hbar}$ |
| <p>(b) It may be regarded as having a wave function that is the superposition of a left-moving particle and a right-moving particle</p> | <p>(f) Valid wave functions include</p> $\Psi(x) = N \cos kx$ <p>where N is a normalization constant and k is defined in (e) above</p> |
| <p>(c) Its energy levels are all negative</p> | <p>(g) All of the above</p> |
| <p>(d) Its energy levels are not quantized</p> | <p>(h) None of the above</p> |

9. Given a particle of mass m in a box of length L having the wave function $\Psi(x) = \sqrt{\frac{2}{L}} \sin\left(\frac{n\pi x}{L}\right)$, what is the energy of the level corresponding to $n = 4$?

- | | |
|--|--|
| <p>(a) Since this wave function is not an eigenfunction of the Hamiltonian the question cannot be answered</p> | <p>(e) $\frac{8\pi^2\hbar^2}{mL^2}$</p> |
| <p>(b) 8 times the energy of the ground state</p> | <p>(f) (c) and (d)</p> |
| <p>(c) $\langle \Psi p_x^2 \Psi \rangle$</p> | <p>(g) (b) and (e)</p> |
| <p>(d) $8\hbar^2 / mL^2$</p> | <p>(h) None of the above</p> |

10. On which of the below functions does the parity operator Π act in the fashion $\Pi[f(x)] = (1)f(x)$?

- | | |
|--------------------------------|---|
| <p>(a) x</p> | <p>(e) Any eigenfunction of the Hamiltonian</p> |
| <p>(b) x^2</p> | <p>(f) (b) and (d)</p> |
| <p>(c) e^{ix}</p> | <p>(g) (a), (b), and (d)</p> |
| <p>(d) $\cos x$</p> | <p>(h) (b), (d), and (e)</p> |

NAME: _____

Short-answer (20 points)

Prove that, given a pair of normalized but *not* orthogonal functions ψ_1 and ψ_2 , the function $\psi_3 = \psi_2 - S\psi_1$ is orthogonal to ψ_1 if S is the overlap integral of ψ_1 and ψ_2 . Is ψ_3 normalized? (Use the back of the page if necessary).

We evaluate

$$\begin{aligned}\langle \psi_1 | \psi_3 \rangle &= \langle \psi_1 | \psi_2 - S\psi_1 \rangle \\ &= \langle \psi_1 | \psi_2 \rangle - S \langle \psi_1 | \psi_1 \rangle \\ &= S - S \cdot 1 \\ &= 0\end{aligned}$$

which proves the orthogonality (we use normalization of ψ_1 in going from line 2 to 3).

With respect to normalization, we evaluate

$$\begin{aligned}\langle \psi_3 | \psi_3 \rangle &= \langle \psi_2 - S\psi_1 | \psi_2 - S\psi_1 \rangle \\ &= \langle \psi_2 | \psi_2 \rangle - 2S \langle \psi_1 | \psi_2 \rangle + S^2 \langle \psi_1 | \psi_1 \rangle \\ &= 1 - 2S \cdot S + S^2 \\ &= 1 - S^2\end{aligned}$$

so ψ_3 is only normalized for the boring case of $S = 0$ (which means that ψ_1 and ψ_2 were orthogonal to begin with since no change to ψ_2 is required to generate ψ_3).

NAME: _____