

A

1. For the diatomic molecule CD, where C has atomic mass 12 and D has atomic mass 2, what is the reduced mass?

- |            |   |
|------------|---|
| (a) 12 / 7 | (e) 10  |
| (b) 7 / 12 | (f) It depends on the vibrational state         |
| (c) 1 / 7  | (g) Cannot be determined from information given |
| (d) 2 / 7  | (h) None of the above                           |

2. Which of the following statements is/are *true* for a given set of QMHO wave functions corresponding to the same harmonic potential  $V$ ?

- |  |   |
|--|---|
| (a) The ground state energy is above the bottom of the potential               | (e) The wave functions are eigenfunctions of the parity operator                |
| (b) The number of nodes is equal to $n+1$ , where $n$ is the energy level      | (f) The selection rule for spectroscopic transitions is $n \rightarrow n \pm 1$ |
| (c) $\langle T \rangle_n = \langle V \rangle_n = (1/2)\langle E \rangle_n$     | (g) (a), (c), (e), and (f)  |
| (d) The wave functions have zero amplitude beyond the classical turning points | (h) All of the above  |

3. Which of the following statements about angular momentum operators and their eigenvalues and eigenfunctions is/are *false*?

- |   |   |
|---|---|
| (a) $L_+ = (L_-)^*$   | (e) $[L_x, L_y] = i\hbar L_z$   |
| (b) $\langle L^2 \rangle = \langle L_z \rangle^2$ whenever $m_l = l$  | (f) $\langle Y_{l,0}   T   Y_{l,0} \rangle > \langle Y_{l',0}   T   Y_{l',0} \rangle$ if $l > l'$ |
| (c) For each value of $l$ there are $2l + 1$ possible values of $m_l$ | (g) (b) and (e)   |
| (d) $L_+ Y_{l,l} = 0$   | (h) All of the above  |

NAME: \_\_\_\_\_

4. What is the eigenvalue of  $L_z$  for  $\Psi$  if the eigenvalue of  $L^2$  for  $\Psi$  is  $25\hbar^2$  and the eigenvalue of  $(L_x^2 + L_y^2)$  for  $\Psi$  is  $9\hbar^2$ ?

- (a) The Heisenberg uncertainty principle dictates that  $\Psi$  cannot be an eigenfunction for  $L_z$  (e)  $\pm 16\hbar$
- (b)  $16\hbar^2$  (f) 0
- (c)  $8i\hbar^2$  (g)  $\pi$
- (d)  $\pm 4\hbar$  (h) None of the above

5. For a diatomic rigid rotator having reduced mass 3 and bond length 2 a.u., which of the following statements is/are *true*?

- (a) The ground-state energy is zero (e) The moment of inertia is 6 a.u.
- (b) The energy separation between the first and second excited states is  $(1/6)$  a.u. (f) (a) and (b)
- (c) The rotational constant  $B$  is  $(1/2)$  a.u. (g) (e) and (f)
- (d) Transition from the ground state to the state  $J = 1$  is forbidden (h) None of the above

6. For a spin-free hydrogenic wave function, which of the below relationships between quantum numbers is/are always true?

- (a)  $n = l > m_l$  (e)  $n = l + m_l$
- (b)  $n > l > m_l$  (f) (b) and (c)
- (c)  $n > l \geq m_l$  (g) (b) and (e)
- (d)  $n > l + m_l$  (h) None of the above

NAME: \_\_\_\_\_

7. Which of the below statements about electron spin is/are true?

- |  |   |
|--|---|
| (a) The spin quantum number comes from including relativity in the electronic Schrödinger equation | (e) Spin-orbit coupling is proportional to the 8th power of the atomic number |
| (b) Spin couples with orbital angular momentum according to $\mathbf{J} = \mathbf{L} + \mathbf{S}$ | (f) (a) and (c)   |
| (c) For a single electron, the only eigenvalues of $S_z$ are $\pm\hbar$                            | (g) (a), (c), and (d)   |
| (d) Stern and Gerlach discovered electron spin by studying the magnetic moments of Au atoms        | (h) All of the above  |

8. An electron of spin  $\alpha$  is in a 4f orbital. Which of the below sets of quantum numbers  $(n, l, m_l, m_s)$  might describe such an electron?

- |                     |                           |
|---------------------|---------------------------|
| (a) (4, 4, 4, 4)    | (e) (4, 4, 3, 1/2)        |
| (b) (4, 3, 2, 1/2)  | (f) (c) and (e)           |
| (c) (4, 3, 0, -1/2) | (g) (b), (c), (d) and (e) |
| (d) (4, 3, 0, 7/2)  | (h) None of the above     |

9. What is the ground-state ionization potential for a one-electron atom having atomic number  $Z$ ?

- |  |   |
|--|---|
| (a) $Z^2$ a.u.   | (e) 1 a.u.  |
| (b) The negative of the energy of the electron in the 1s orbital | (f) The energy required to infinitely separate the nucleus and electron |
| (c) $(1/2)Z^2$ a.u.  | (g) (b), and (d)  |
| (d) $2Z^2$ a.u.  | (h) (b), (c), and (f)   |

NAME: \_\_\_\_\_

10. Which of the following wave functions has the greatest degeneracy?

- |  |  |
|--|--|
| (a) Particle in a box, level $n = 8$                       | (e) Spin-free hydrogenic wave function, $n = 3$                                  |
| (b) Rigid rotator, $l = 4$                                 | (f) Relativistic free electron at rest   |
| (c) Quantum mechanical harmonic oscillator, level $n = 25$ | (g) (b) and (e) have the same degeneracy which is greater than all of the others |
| (d) Spin-free hydrogenic wave function, $n = 6, l = 1$     | (h) (a) through (f) are all singly degenerate                                    |

**Short answer.** Show that by proper choice of  $a$ , the function  $e^{-ar^2}$  is an eigenfunction of the operator

$$\left[ \frac{d^2}{dr^2} - qr^2 \right]$$

where  $q$  is a constant. What is the name of the general class of functions represented by  $e^{-ar^2}$ ? How many nodes does this function have over  $r$ ?

To show that  $e^{-ar^2}$  is an eigenfunction with proper choice of  $a$  we require

$$\left[ \frac{d^2}{dr^2} - qr^2 \right] e^{-ar^2} = ze^{-ar^2}$$

Evaluating the l.h.s. we have

$$\begin{aligned} \left[ \frac{d^2}{dr^2} - qr^2 \right] e^{-ar^2} &= -2ae^{-ar^2} + 4a^2r^2e^{-ar^2} - qr^2e^{-ar^2} \\ &= -\left[ 2a + (q - 4a^2)r^2 \right] e^{-ar^2} \end{aligned}$$

and for the prefactor on the r.h.s. to be a constant (so as to satisfy the eigenvalue condition) it must be true that  $a$  is  $\sqrt{q}/2$  in which case the eigenvalue will be  $-2a$ , which is simply  $-\sqrt{q}$ .

The general class of functions here are “gaussian” functions. A gaussian has no nodes over  $r$ .

NAME: \_\_\_\_\_